

球面上の超対称ゲージ理論の数値実験

Numerical Experiment of Supersymmetric Gauge Theory on 2-Sphere

So Matsuura

(Department of Physics Hiyoshi, Keio University)

Based on work with K. Ohta, T. Misumi and S. Kamata

Continuum theory

Starting point 2D $N = (2,2)$ SYM theory

bosons: $A_\mu, \phi, \bar{\phi}$

fermions: $\Psi = \begin{pmatrix} \bar{\psi} \\ \psi \end{pmatrix}$

$$S_{2d} = \frac{1}{2g^2} \int d^2x \text{Tr} \left\{ \frac{1}{2} F_{\mu\nu}^2 + (\mathcal{D}_\mu \phi) (\mathcal{D}^\mu \bar{\phi}) + \frac{1}{4} [\phi, \bar{\phi}]^2 \right. \\ \left. + i \bar{\Psi} \Gamma^\mu \mathcal{D}_\mu \Psi + \frac{i}{2} \bar{\Psi} \Gamma_+ [\bar{\phi}, \Psi] + \frac{i}{2} \bar{\Psi} \Gamma_- [\phi, \Psi] \right\}.$$

global symmetries

$U(1)_R$ transformation

$$A_\mu \rightarrow A_\mu, \quad \phi \rightarrow e^{2i\theta} \phi, \quad \bar{\phi} \rightarrow e^{-2i\theta} \bar{\phi}, \quad \psi \rightarrow e^{i\theta} \psi, \quad \bar{\psi} \rightarrow e^{-i\theta} \bar{\psi}.$$

$U(1)_V$ transformation


$$A_\mu \rightarrow A_\mu, \quad \phi \rightarrow \phi, \quad \bar{\phi} \rightarrow \bar{\phi}, \quad \psi \rightarrow e^{i\theta\sigma_2} \psi, \quad \bar{\psi} \rightarrow e^{i\theta\sigma_2} \bar{\psi}.$$

SUSY transformation

$$\delta A_\mu = \bar{\epsilon}^T \gamma_\mu \psi + \epsilon^T \gamma_\mu \bar{\psi}, \quad \delta \phi = -2i \bar{\epsilon}^T \bar{\psi}, \quad \delta \bar{\phi} = 2i \epsilon^T \psi, \\ \delta \psi = -i F_{12} \sigma_2 \epsilon - i (\mathcal{D}_\mu \bar{\phi}) \gamma_\mu \bar{\epsilon} + \frac{i}{2} [\phi, \bar{\phi}] \epsilon, \quad \delta \bar{\psi} = -i F_{12} \sigma_2 \bar{\epsilon} + i (\mathcal{D}_\mu \phi) \gamma_\mu \epsilon - \frac{i}{2} [\phi, \bar{\phi}] \bar{\epsilon},$$


Derivatives on a curved background

$$\begin{aligned}\partial_\mu &\rightarrow \nabla_\mu && \text{spin connection: } \omega_{12} \equiv \omega \\ \nabla_\mu \psi &= (\partial_\mu + \frac{1}{4} \omega_\mu^{ab} \gamma_{ab}) \psi = \partial_\mu \psi - \frac{i}{2} \omega_\mu \sigma_2 \psi, \\ \nabla_\mu \bar{\psi} &= (\partial_\mu - \frac{1}{4} \omega_\mu^{ab} \gamma_{ab}) \bar{\psi} = \partial_\mu \bar{\psi} + \frac{i}{2} \omega_\mu \sigma_2 \bar{\psi}.\end{aligned}$$



Derivatives on a curved background + background $U(1)_V$ field B_μ

$$\begin{aligned}\partial_\mu \psi &\rightarrow (\nabla_\mu + \frac{i}{2} B_\mu \sigma_2) \psi = \left(\partial_\mu + \frac{i}{2} (B_\mu - \omega_\mu) \sigma_2 \right) \psi, \\ \partial_\mu \bar{\psi} &\rightarrow (\nabla_\mu + \frac{i}{2} B_\mu \sigma_2) \bar{\psi} = \left(\partial_\mu + \frac{i}{2} (B_\mu + \omega_\mu) \sigma_2 \right) \bar{\psi},\end{aligned}$$



Special background or Topological twisting [Seiberg 2011, 2012](#)

$$B_\mu = \omega_\mu \quad \longrightarrow \quad \begin{aligned} \left(\partial_\mu + \frac{i}{2} (B_\mu - \omega_\mu) \sigma_2 \right) \psi &\rightarrow \partial_\mu \psi && \text{: scalar} \\ \left(\partial_\mu + \frac{i}{2} (B_\mu + \omega_\mu) \sigma_2 \right) \bar{\psi} &\rightarrow (\partial_\mu + i \omega_\mu \sigma_2) \bar{\psi} && \text{: vector} \end{aligned}$$

Natural renaming in this background

$$\psi \rightarrow (\chi, \eta/2)^T : \text{scalar}$$

$$\bar{\psi} \rightarrow \lambda_\mu : \text{vector}$$

Continuum action

$$S_{2d} = \frac{1}{2g^2} \int d^2x \sqrt{g} \text{Tr} \left\{ (F_{12}/\sqrt{g})^2 + (\mathcal{D}_\mu \phi) (\mathcal{D}^\mu \bar{\phi}) + \frac{1}{4} [\phi, \bar{\phi}]^2 \right. \\ \left. - 2i E^{\mu\nu} \lambda_\mu \mathcal{D}_\nu \chi + i \lambda^\mu \mathcal{D}_\mu \eta + \lambda^\mu [\bar{\phi}, \lambda_\mu] - \psi^a [\phi, \psi_a] \right\}, \\ (E^{\mu\nu} = \varepsilon^{\mu\nu} / \sqrt{g})$$

$U(1)_R$ symmetry

$$\delta A_\mu = 0, \quad \delta \Phi = 2i\theta \Phi, \quad \delta \bar{\Phi} = -2i\theta \bar{\Phi} \\ \delta \lambda_\mu = i\theta \lambda_\mu, \quad \delta \eta = -i\theta \eta, \quad \delta \chi = -i\theta \chi.$$

$U(1)_V$ symmetry

$$\delta \psi_a = \theta \varepsilon_{ab} \psi_b, \\ \delta \lambda_\mu = -\theta E_{\mu\nu} \lambda^\nu.$$

SUSY transformation of the action

(1) scalar SUSY transformations: $\epsilon_a = \epsilon_a(x) \quad \bar{\epsilon}_\mu = 0 \quad \left(C^{1;\mu\nu} = \frac{-1}{\sqrt{g}} \epsilon^{\mu\nu}, \quad C^{2;\mu\nu} = g^{\mu\nu} \right)$

$$\delta S = \frac{1}{2g^2} \int d^2x \sqrt{g} (\partial_\mu \epsilon^a) J_a^\mu(x) \quad \text{preserve for } \epsilon_a = \text{const.}$$

$$\left(J_a^\mu(x) = 2i \text{Tr} \left\{ E^{\mu\nu} (\lambda C_a)_\nu f + (\mathcal{D}^\mu \phi) \psi_a + \varepsilon_{ab} E^{\rho\nu} (\mathcal{D}_\rho \phi) \psi_b - \frac{i}{2} (\lambda C_a)^\mu [\phi, \bar{\phi}] \right\} \right)$$

(2) vector SUSY transformations: $\epsilon_a = 0 \quad \bar{\epsilon}_\mu = \bar{\epsilon}_\mu(x)$

$$\bar{\delta} S = \frac{1}{2g^2} \int d^2x \sqrt{g} (\nabla^\mu \bar{\epsilon}^\nu) \bar{J}_{\mu\nu}(x) \quad \text{preserve iff } \bar{\epsilon}_\mu \text{ is covariantly const.}$$

$$\left(\bar{J}_{\mu\nu}(x) = 2i \text{Tr} \left\{ \frac{i}{2} C_{a;\nu\mu} \psi_a [\phi, \bar{\phi}] - \sigma_{ab}^1 C_{b;\mu\nu} \psi_a f - \lambda_\nu \mathcal{D}_\mu \bar{\phi} - \lambda_\mu \mathcal{D}_\nu \bar{\phi} + g_{\mu\nu} \lambda^\rho \mathcal{D}_\rho \bar{\phi} \right\} \right)$$

$$\delta = i\epsilon_1 \tilde{Q} + i\epsilon_2 Q$$

$$\left\{ \begin{array}{ll} QA_\mu = \lambda_\mu, & Q\lambda_\mu = i\mathcal{D}_\mu\phi, \\ Q\bar{\phi} = \eta, & Q\eta = [\phi, \bar{\phi}], \\ Q\chi = Y, & QY = [\phi, \chi], \\ Q\phi = 0 \end{array} \right. \quad \left\{ \begin{array}{ll} \tilde{Q}A_\mu = \sqrt{g}\varepsilon_{\mu\nu}\lambda^\nu, & \tilde{Q}\lambda_\mu = -i\sqrt{g}\varepsilon_{\mu\nu}\mathcal{D}^\nu\phi, \\ \tilde{Q}\bar{\phi} = 2\chi, & \tilde{Q}\chi = \frac{1}{2}[\phi, \bar{\phi}], \\ \tilde{Q}\eta = -2Y, & \tilde{Q}Y = -\frac{1}{2}[\phi, \eta], \\ \tilde{Q}\phi = 0 \end{array} \right.$$

Continuum action in Q-exact form

$$S_0 = \frac{1}{2g^2} \int d^2x \sqrt{g} Q \text{Tr} \left\{ \frac{1}{4} \eta[\phi, \bar{\phi}] - ig^{\mu\nu} \lambda_\mu \mathcal{D}_\nu \bar{\phi} + \chi \left(Y - \frac{2i}{\sqrt{g}} F_{12} \right) \right\}$$



Observation toward discretized theory

Bosonic fields on lattice

scalar $\phi(x)$ $\longrightarrow$ site variable

vector $A_\mu(x)$ $\longrightarrow$ link variable

field tensor $F_{\mu\nu}(x)$ $\longrightarrow$ face variable




requirement

(1) assign bosons on the lattice corresponding to their vector structure

(2) keep Q-symmetry

Fields on lattice



Discretized action cf) Sugino 2003

$$S = S_S + S_L + S_F \equiv Q \left\{ \sum_{s \in S} \alpha_s \Xi_s + \sum_{\langle st \rangle \in L} \alpha_{\langle st \rangle} \Xi_{\langle st \rangle} + \sum_{f \in F} \alpha_f \Xi_f \right\}$$

$$\left\{ \begin{array}{l} \Xi_s \equiv \frac{1}{2g_0^2} \text{Tr} \left\{ \frac{1}{4} \eta_s [\Phi_s, \bar{\Phi}_s] \right\}, \\ \Xi_{\langle st \rangle} \equiv \frac{1}{2g_0^2} \text{Tr} \left\{ -i \lambda_{st} (U_{st} \bar{\Phi}_t U_{st}^{-1} - \bar{\Phi}_s) \right\}, \\ \Xi_f \equiv \frac{1}{2g_0^2} \text{Tr} \left\{ \chi_f (Y_f - i \beta_f \mu(U_f)) \right\}, \end{array} \right.$$

$$\left\{ \begin{array}{ll} Q\Phi_s = 0, & Q\eta_s = [\Phi_s, \bar{\Phi}_s], \\ Q\bar{\Phi}_s = \eta_s, & Q\lambda_{st} = i (U_{st} \Phi_t U_{st}^{-1} - \Phi_s + \lambda_{st} \lambda_{st}), \\ QU_{st} = i \lambda_{st} U_{st}, & Q\chi_f = Y_f, \\ QY_f = [\Phi_f, \chi_f], & \end{array} \right.$$

$$\left(U_f \equiv \prod_{i=1}^n U_{s_i s_{i+1}} \right)$$

Topological information is preserved on lattice

(1) The same localization technique with the continuum theory works also on lattice

non-trivial fixed point equations

$$\begin{aligned}\Phi_s U_{st} - U_{st} \Phi_t &= 0 \\ \mu(U_f) &= 0\end{aligned}$$



1-loop contribution is exact

$$\begin{aligned}Z &\sim \frac{(\Delta(\phi)^2)^{\#(\text{sites})} \times (\Delta(\phi))^{\#(\text{faces})}}{(\Delta(\phi))^{\#(\text{links})} \times (\Delta(\phi))^{\#(\text{sites})}} \\ &= \prod_{i < j} (\phi_i - \phi_j)^\chi \quad \leftarrow \text{Euler characteristic !}\end{aligned}$$

(2) The $U(1)_R$ anomaly appears in the measure

$$\mathcal{D}\vec{B} = \left(\prod_{s=1}^{N_S} \mathcal{D}\Phi_s \mathcal{D}\bar{\Phi}_s \right) \left(\prod_{l=1}^{N_L} \mathcal{D}U_l \right) \left(\prod_{f=1}^{N_F} \mathcal{D}Y_f \right) : U(1)_R \text{ neutral}$$

$$\mathcal{D}\vec{F} = \left(\prod_{s=1}^{N_S} \mathcal{D}\eta_s \right) \left(\prod_{l=1}^{N_L} \mathcal{D}\lambda_l \right) \left(\prod_{f=1}^{N_F} \mathcal{D}\chi_f \right)$$

$$\rightarrow \left(\mathcal{D}\vec{\mathcal{F}} \right) e^{i(N_S - N_L + N_F)(N^2 - 1)\alpha}$$

$U(1)_R$ symmetry

$$\begin{aligned}\Phi &\rightarrow e^{2i\alpha} \Phi, & \bar{\Phi} &\rightarrow e^{-2i\alpha} \bar{\Phi}, & A_\mu &\rightarrow A_\mu, \\ \eta &\rightarrow e^{-i\alpha} \eta, & \lambda_\mu &\rightarrow e^{i\alpha} \lambda_\mu, & \chi &\rightarrow e^{-i\alpha} \chi.\end{aligned}$$

Natural question : Can we take the continuum limit ?

YES! from power counting point of view

(1) Tree level continuum limit reproduces the continuum action:

$$S = S_S + S_L + S_F \equiv Q \left\{ \sum_{s \in S} \alpha_s \Xi_s + \sum_{\langle st \rangle \in L} \alpha_{\langle st \rangle} \Xi_{\langle st \rangle} + \sum_{f \in F} \alpha_f \Xi_f \right\}$$

$$\begin{aligned} a^2 &\equiv \text{vol}(\Sigma_g)/N_F & a^2 A_f &= \int_{\sigma_f} d^2x \sqrt{g(x)}, & e_{st}^\mu &\equiv \frac{1}{a} (x_t^\mu - x_s^\mu) \\ \alpha_s &= \sum_{f \in F_s} \frac{A_f}{|S_f|}, & \alpha_f &= A_f, & \beta_f &= \frac{1}{A_f}, & \sum_{\langle st \rangle \in L_f} \alpha_{\langle st \rangle}^f e_{st}^\mu e_{st}^\nu &= A_f g^{\mu\nu}(x_f). \end{aligned}$$

(2) There is no **Q**, **$U(1)_R$** and **gauge-invariant** radiative correction which spoils the geometry.

$$\left(\underbrace{\frac{a^{p-4}}{g^2}}_{\text{tree}} + \underbrace{c_1 p^{p-2}}_{\text{1-loop}} + \underbrace{c_2 a^p g^2}_{\text{2-loop}} + \cdots \right) \int d^2x \sqrt{g} \mathcal{O}_p(x) \quad \longrightarrow \quad \mathcal{O} \sim \mathcal{B} \text{ or } \mathcal{B}^2$$

It is expected that the continuum theory will be obtained by simply taking $a \rightarrow 0$.

We have to check it non-perturbatively.

(It gives a “definition” of the SUSY theory on curved background!)

What should we check?

The $U(1)_V$ symmetry and $\tilde{Q}$ -symmetry are broken by discretization.

➡ They must be restored in the continuum limit.

fact These symmetries are related with each other in the continuum limit.

$$J_V^\mu = 2i \text{Tr} \left\{ \varepsilon_{ab} (\lambda C_a)_\mu \psi_b \right\}.$$

$$J_a^\mu(x) = 2i \text{Tr} \left\{ E^{\mu\nu} (\lambda C_a)_\nu f + (\mathcal{D}^\mu \phi) \psi_a + \varepsilon_{ab} E^{\rho\nu} (\mathcal{D}_\rho \phi) \psi_b - \frac{i}{2} (\lambda C_a)^\mu [\phi, \bar{\phi}] \right\}$$

$$Q_a J_V^\mu(x) = \varepsilon_{ab} J_b^\mu(x)$$

$$dJ_Q = 0$$

$$dJ_V = 0 \quad \longrightarrow \quad dJ_{\tilde{Q}} = 0$$

It is sufficient to check the $U(1)_V$ symmetry

Model

(M,N)-polygon decomposition of S^2

parameter tuning

$$ds^2 = R^2(d\theta^2 + \cos^2 \theta d\varphi^2) \quad (-\pi/2 < \theta < \pi/2)$$

$$a^2 = \frac{8\pi R^2}{M(N-2) + 4}$$

$$\alpha_{s=1\dots M} = \alpha_{s=\frac{MN}{2}-M+1\dots\frac{MN}{2}} = \frac{\pi R^2}{Ma^2} \left(2 - \cos \frac{\pi}{N} - \cos \frac{3\pi}{N} \right)$$


$$\alpha_{s=nM+i} = \frac{\pi R^2}{Ma^2} \left(\cos \frac{2n-1}{N} \pi - \cos \frac{2n+3}{N} \pi \right)$$

$$\alpha_{l=(n-1)M+i} = \frac{M}{\pi} \sin \frac{\pi}{N} \left(\sum_{k=1}^{n-1} - \sum_{k=n}^{N/2-1} \right) \frac{(-1)^{n+k+1}}{\sin \frac{2k\pi}{N}}$$

$$\alpha_{l=MN/2+(n-1)M+i} = \frac{N^2}{2\pi M} \left(\cos \frac{2n-1}{N} \pi - \cos \frac{2n+1}{N} \pi \right)$$

$$\alpha_{f=(n-1)M+i} = \beta_{f=(n-1)M+i}^{-1} = \frac{2\pi R^2}{a^2} \left(\cos \frac{2n-1}{N} \pi - \cos \frac{2n+1}{N} \pi \right)$$

$$\alpha_{f=1, M(N-2)/2+2} = \beta_{f=1, M(N-2)/2+2}^{-1} = \frac{2\pi R^2}{a^2} \left(1 - \cos \frac{\pi}{N} \right)$$



site	$N \times M/2$
link	$N \times M/2 + M \times (N-2)/2$
face	$2 + M \times (N-2)/2$

Anomaly-phase-quench method

vev in the continuum theory

$$\langle \mathcal{O} \rangle \equiv \frac{1}{Z_q} \int \mathcal{D}\vec{B} \mathcal{D}\vec{F} \mathcal{O} e^{-S_b - S_f} = \frac{1}{Z_q} \int \mathcal{D}\vec{B} \mathcal{O} \text{Pf}(D) e^{-S_b}$$

U(1) charge: $(N^2 - 1)\chi_h$

phase quench method is usually used in Monte Carlo method

$$\langle \mathcal{O} \rangle^q \equiv \frac{1}{Z_q} \int \mathcal{D}\vec{B}' \mathcal{O} |\text{Pf}(D)| e^{-S_b}$$

U(1) charge: ZERO

**NOT A GOOD
APPROXIMATION**

philosophy of the phase quench

Ignore only the artificial phase
coming from the discretization

Observation

$$\text{Pf}(D) = |\text{Pf}(D)| e^{i\theta_A + i\theta}$$

1. U(1)_R phase θ_A

2. lattice artifact θ

We should ignore only θ

Compensator

Kamata-Misumi-Ohta-S.M. 2016

$\mathcal{A}$: an operator with

- $Q\mathcal{A} = 0$
- $[\mathcal{A}] = -(N^2 - 1)\chi_h$
- $\mathcal{A} \equiv |\mathcal{A}|e^{-i\theta_A}$

anomaly-phase-quench method

$$\langle \mathcal{O} \rangle^{\hat{q}} \equiv \langle \mathcal{O} e^{i\theta_A} \rangle^q = \frac{1}{Z_q} \int \mathcal{D}\vec{B} \mathcal{O} |Pf(\mathcal{D})| e^{i\theta_A}$$

trace type

$$\mathcal{A}_{\text{tr}} = \frac{1}{N_S} \sum_{s=1}^{N_S} \left(\frac{1}{N_c} \text{Tr}(\Phi_s)^2 \right)^{-\frac{N_c^2-1}{4}\chi_h}$$

determinant type

$$\mathcal{A}_{\text{det}} = \frac{1}{N_S} \sum_{s=1}^{N_S} (\text{Det}\Phi_s)^{-\frac{N_c^2-1}{2N_c}\chi_h}$$

Izykson-Zuber type



$$\mathcal{A}_{\text{IZ}} = \frac{1}{N_l} \sum_{l=1}^{N_l} \left(\frac{1}{N_c} \text{Tr} \left(2\Phi_{\text{org}(l)} U_l \Phi_{\text{tip}(l)} U_l^\dagger + \lambda_l \lambda_l (U_l \Phi_{\text{tip}(l)} U_l^\dagger + \Phi_{\text{org}(l)}) \right) \right)^{-\frac{N_c^2-1}{4}\chi_h}$$

Trivial WT identity for Q -symmetry

basic relation (for $\mu = 0$)

$$0 = \langle Q\Xi \rangle = \langle S_b + S_f \rangle = \langle S_b \rangle - \frac{\#(\text{fermions})}{2}$$

$$\longrightarrow \langle \mathcal{A} S_b \rangle + \frac{\mu^2}{2} \langle \mathcal{A} \Xi \sum_s \text{Tr}(\Phi_s \eta_s) \rangle = \frac{N_c^2 - 1}{2} (N_S + N_L) \langle \mathcal{A} \rangle$$



Identity 1 from $U(1)_V$ symmetry

Site action

$$\frac{1}{2g_0^2} \sum_s \alpha_s Q \text{Tr} \left(\frac{1}{4} \eta_s [\phi_s, \bar{\phi}_s] \right) = \frac{1}{2g_0^2} \sum_s \alpha_s \text{Tr} \left(\frac{1}{4} [\phi_s, \bar{\phi}_s]^2 \right) - \frac{1}{2g_0^2} \sum_s \alpha_s \text{Tr} \left(-\frac{1}{4} \eta_s [\phi_s, \eta_s] \right)$$

Face action

Q -exact


$$\frac{1}{2g_0^2} \sum_f \alpha_f Q \text{Tr} (\chi_f Y_f) = \frac{1}{2g_0^2} \sum_f \alpha_f \text{Tr} (Y_f^2) - \frac{1}{2g_0^2} \sum_f \alpha_f \text{Tr} (-\chi_f [\phi_f, \chi_f])$$

$U(1)_V$ doublet



$$\left\langle \frac{1}{2g_0^2} \sum_s \alpha_s \text{Tr} \left(\frac{1}{4} [\phi_s, \bar{\phi}_s]^2 \right) \right\rangle + \left\langle \frac{1}{2g_0^2} \sum_f \frac{\alpha_f \beta_f^2}{4} \text{Tr} (\Omega(U_f)^2) \right\rangle$$

$$\rightarrow \frac{1}{2} N_F (N_c^2 - 1) \quad (a \rightarrow 0)$$




Identity 2 from $U(1)_V$ symmetry

$$\frac{1}{2g_0^2} \sum_l \alpha_l Q \text{Tr} (-i\lambda_l (U_l \bar{\phi}_{t(l)} U_l^{-1} - \bar{\phi}_{s(l)})) = \frac{1}{2g_0^2} \sum_l \alpha_l \text{Tr} (|D_l \phi|^2 + i\lambda_l D_l \eta - \lambda_l \lambda_l (U_l \bar{\phi}_{t(l)} U_l^{-1} + \bar{\phi}_{s(l)}))$$


Q -exact

$$\frac{1}{2g_0^2} \sum_f \alpha_f Q \text{Tr} (-i\beta_f \chi_f \Omega(U_f)) = \frac{1}{2g_0^2} \sum_f \alpha_f \beta_f \text{Tr} (-iY_f \Omega(U_f) + i\chi_f Q \Omega(U_f))$$

$U(1)_V$ doublet



$$2\langle S_B^F \rangle + \left\langle \frac{1}{2g_0^2} \sum_l \alpha_l \text{Tr} (\lambda_l \lambda_l (U_l \bar{\phi}_{t(l)} U_l^{-1} + \bar{\phi}_{s(l)})) \right\rangle \rightarrow \langle S_B^L \rangle \quad (a \rightarrow 0)$$



We should check $U(1)_V$ WT identity

In the continuum theory

$$J_V^\mu = \text{Tr} \left\{ \frac{1}{2} E^{\mu\nu} \lambda_\nu \eta - \lambda^\mu \chi \right\}$$

For a $U(1)_V$ invariant operator $\mathcal{O}(x)$,


$$\left\langle \mathcal{O}(y) \mathcal{D}^\mu J_\mu^V(x) \right\rangle = \left\langle \mathcal{O}(y) \left(\frac{1}{2} \nabla \times \text{Tr}(\lambda \eta) - \nabla \cdot \text{Tr}(\lambda \chi) \right) \right\rangle = 0$$

problem 1

There are infinitely many ways to construct **composite operators** on the lattice.

problem 2

How to define **Rotation** and **Divergence** on lattice?



Hint: exact $U(1)_R$ current on the lattice

cf) Ohta-san's talk

continuum

$$\delta S_{cont} = \frac{1}{2g^2} \int d^2x \sqrt{g} (2i\partial_\mu \theta) \text{Tr} \left\{ 2i(\phi \mathcal{D}^\mu \bar{\phi} - \bar{\phi} \mathcal{D}^\mu \phi) + 2E^{\mu\nu} \lambda_\nu \chi + \lambda^\mu \eta \right\}$$

lattice ($\theta_s \equiv \theta \delta_{s,s_0}$)

$$\begin{aligned} \delta S_{lat} = & \frac{2i\theta}{2g_0^2} \left(\sum_{l \in \langle \cdot, s_0 \rangle} - \sum_{l \in \langle s_0, \cdot \rangle} \right) \alpha_l \text{Tr} (\Phi_{s(l)} U_l \bar{\Phi}_{t(l)} U_l^{-1} - \bar{\Phi}_{s(l)} U_l \Phi_{t(l)} U_l^{-1}) \\ & \text{DIVERGENCE} \\ & + \frac{2i\theta}{2g_0^2} \left(\sum_{l \in \langle \cdot, s_0 \rangle} - \sum_{l \in \langle s_0, \cdot \rangle} \right) \alpha_l \text{Tr} \left(\frac{i}{2} \lambda_l U_l \eta_{t(l)} U_l^{-1} + \lambda_l \lambda_l U_l \bar{\Phi}_{t(l)} U_l^{-1} \right) \\ & \text{DIVERGENCE} \\ & + \frac{2i\theta}{2g_0^2} \left(\sum_{l \in \langle s_0, \cdot \rangle} \sum_{f \in F_l} - \sum_{f=s_0} \sum_{l \in L_f} \right) (i\alpha_f \beta_f \epsilon_{f,l}) \left\{ \frac{1}{2B(U_f)} \text{Tr} (\chi_f X_{f,l} \lambda_l Y_{f,l} + \chi_f Y_{f,l}^\dagger \lambda_l X_{f,l}^\dagger) \right. \\ & \text{ROTATION} \\ & \left. - \frac{1}{\epsilon^2} \frac{1}{2B(U_f)^2} \text{Tr} (\chi_f (U_f - U_f^{-1})) \text{Tr} (X_{f,l} \lambda_l Y_{f,l} - Y_{f,l}^\dagger \lambda_l X_{f,l}^\dagger) \right\} \end{aligned}$$

$B(U_f) = 1 - \frac{1}{\epsilon^2} \text{Tr} (2 - U_f - U_f^{-1}).$

This correspondence will be a guiding principle to construct
 “ $U(1)_V$ current”, rotation and divergence on the lattice.

on going...

Summary and Future Works

- We explicitly discretize 2-sphere.
- Q-symmetry is preserved as expected.
- The relations expected from the $U(1)_V$ symmetry seems to hold in the continuum limit.
- $U(1)_V$ WT identity, restoration of S^2 geometry Kamata-Ohta-Misumi-S.M. to be appeared
- Other discretization (fullerene-like discretization?)
- $N=(4,4)$ and $N=(8,8)$ theory
- Simulation of 4D $N=4$ SYM by hybrid method cf) Hanada-Sugino-S.M. 2012
- Theory with matters
- Connection to Dynamical triangulation? (supersymmetric dynamical triangulation?)